



Great Northern
LANDSCAPE CONSERVATION COOPERATIVE



Strategic Conservation Framework

Contents

Introduction	1
Vision	2
Organizational Structure	3
Scope	4
The Great Northern Landscape	7
Conservation Goals	8
Conservation Targets	10
Columbia Basin	12
Rocky Mountains	14
Sagebrush Steppe	16
Next Steps	18
Acknowledgements	19

Suggested citation:

Chambers, N., G. Tabor, Y. Converse, T. Olliff, S. Finn, R. Sojda, and S. Bischke. 2013. The Great Northern Landscape Conservation Cooperative Strategic Conservation Framework.

Cover photos (photo credit): Morning light on King Edward Peak, British Columbia (Steve Gnam), Canadian lynx (NPS), cutthroat trout (USGS), mule deer (NPS), aspen (Montana Department of Environmental Quality), Columbia Gorge (Northwest Council), sagebrush (NPS)

Foreword

The purpose of this strategic conservation framework is to articulate the rationale, approach, and priorities for the Great Northern Landscape Conservation Cooperative (GNLCC) that reflects the unique geography and regional natural resource issues. The information presented in this document is summarized from background research on existing landscape initiatives (place-, issue-, or species-based) and other regionally summarized ecological and landscape information relevant to the Great Northern geography.

The conservation targets identified in this document are based on research with conservation partners and ongoing landscape-scaled initiatives. We collected and reviewed over 50 documents that may be relevant to the Great Northern LCC and conducted nearly 60 interviews with members of the Steering Committee, Advisory Team, and Partner Forum representatives. The conservation targets (i.e., ecosystem processes, ecosystems or habitats, and species or focal resources) provide a common focus for organizations to move toward a collective landscape vision using specific ecological outcomes within the GNLCC.

The GNLCC Strategic Conservation Framework (as described in the GNLCC Governance Charter) is intended to guide priority setting for annual work plans over a five- to ten-year time period. The framework provides the foundation for specific research and technical actions, institutional collaboration, and the application of ecological information, including future additions through GNLCC Partner Forums (see the GNLCC Governance Charter).

We recognize that this document is a starting point, and that the priorities expressed herein will continue to evolve and be refined. We are grateful to the advisory team and partners who have helped shape this document, and to our colleagues leading the partner forums, who will take the concepts in this framework the next steps.

Yvette Converse
Tom Olliff
Great Northern LCC Coordinators



PHOTO CREDIT: NPS, Nancy Bookino

Introduction

Landscape Conservation Cooperatives (LCCs) are applied conservation science partnerships that provide scientific and technical support for conservation and inform sustainable resource management at landscape scales. Landscape Conservation Cooperatives are intended to inform sustainable management of land, water, fish, wildlife, and cultural heritage resources in response to climate change and other landscape-level challenges. The Great Northern LCC (GNLCC) is one of 22 LCCs in an international network established by U.S. Department of the Interior Secretarial Order 3289. For a map and more information about LCCs, go to <http://www.doi.gov/lcc/index.cfm>.

The GNLCC is a voluntary network of partners working to address common landscape conservation goals. We work together to identify commonalities among our efforts and build consensus to enact a regional approach to landscape conservation. We work across boundaries and jurisdictions, and share data, science, and capacity to achieve common goals.

The GNLCC is creating the conditions that enhance individual and collective partner implementation of landscape-level conservation. Through information sharing, capacity building, effective analyses and decision-support tools, and collaborative networks, the GNLCC creates efficiencies and reduces the challenges of working in complex ecological and jurisdictional systems.

The GNLCC supports the missions and mandates of its partners. Working together in partnership, the GNLCC can add value to existing efforts through:

- Creating a vehicle for collaborative, collective effort to address landscape-level challenges;
- Putting science in the hands of practitioners through shared information, accessible tools, and resources; and
- Facilitating the interaction of existing partnerships and initiatives and supporting alignment for greater effectiveness.

Vision

A landscape that sustains its diverse natural systems to support healthy and connected populations of fish, wildlife, and plants; sustains traditional land uses and cultural history; and supports robust communities.

Mission

Coordinate, facilitate, promote, and add value to large landscape conservation to build resource resilience and inform sustainable resource management in the face of climate change and other landscape stressors through the following:

SUPPORT SCIENCE DEVELOPMENT

Identify and facilitate the development, integration, and application of social and natural scientific information needed to inform conservation actions and sustainable resource management decisions to conserve water, land, fish, wildlife, and cultural heritage.

EFFECT COORDINATION

Support coordination and integration of conservation science and actions across ecosystems at the landscape scale, leveraging the capabilities of respective agencies/organizations/partnerships, and provide real-time situational awareness of on-going conservation efforts.

INFORM CONSERVATION ACTION AND SUSTAINABLE RESOURCE MANAGEMENT

Provide scientific information and decision support tools informing conservation action and sustainable resource management that conserve water, land, fish, wildlife, and cultural heritage.

MONITOR AND EVALUATE

Support efforts to monitor landscape-scale indicators, test scientific assumptions, and evaluate effectiveness of conservation actions and sustainable resource management to inform adaptive management decision making.

COMMUNICATE AND EDUCATE

Communicate relevant science information and GNLCC activities and opportunities to partners and users. Facilitate collaborative conservation and seek to leverage capabilities and support.

Organizational Structure

The Great Northern LCC is a network of partners working toward a collective landscape vision and conservation goals. This network comprises diverse U.S. federal, Canadian provincial and federal,

Tribal Nations, state, nongovernmental, and academic conservation organizations. These groups work together within an organizational structure as described in Figure 1.



Figure 1. Organizational structure of the GNLCC, identifying main committees, their roles, and functions.



Scope

The Great Northern is an international, transboundary landscape and a complex mosaic of geography, organizations, and jurisdictions. The U.S. portion includes large parts of Idaho, Montana, Oregon, Washington, and Wyoming—an area of approximately 260 million acres. A large portion of British Columbia (BC), and a portion of Alberta, Canada make up approximately 40 percent of the Great Northern region. The GNLCC has embraced an international approach to transboundary conservation; however, there are fundamental differences in government structure and jurisdiction

among the partners. Canadian partners are coordinating to determine the right model of landscape conservation that can interface with U.S. efforts within the Great Northern area. Collectively, this will define the GNLCC; at this point, however, this strategy was developed primarily through coordination of U.S. partners with involvement from Canadian partners in transboundary issues. It is recognized that this strategy may be broadened at some point in the future to represent the priorities of the Canadian partners, as the organizational model and appropriate international interface is developed.



PHOTO CREDIT: Gary Tabor

Guiding Principles

WORK COOPERATIVELY AND COLLABORATIVELY to improve effectiveness of each organization's large-scale landscape conservation programs and efforts.

CONDUCT OPEN AND FREQUENT COMMUNICATIONS within the GNLCC network, between related climate change and landscape programs, and among the expanded climate change and landscape conservation community.

CONSIDER AND RESPECT each participating organization's unique mandate and jurisdiction.

COORDINATE WITH OTHER committees, work groups, or organizations that add mutual value, maximize capacity, avoid redundancies, and leverage resources.

FOCUS ON SOLVING scientific, ecological, and biological issues to promote scientifically sound, outcome-based adaptive management.

RESPECT SOCIAL, POLITICAL, AND LEGAL LIMITATIONS while promoting solutions to landscape-level stressors that benefit the greater GNLCC conservation community.

BE TRANSPARENT in operations and ensure equal and open access.




Figure 2. Map of land cover in the GNLCC landscape.

The Great Northern Landscape

As a landscape, the Great Northern area includes interior British Columbia and portions of Alberta in Canada, the eastern portion of Washington, northeastern Oregon, northern Idaho, and western Montana and Wyoming. It includes a large portion of the Rocky Mountains (central and north) as well as portions of two major U.S. river basins (the Upper Columbia and Upper Missouri) with topography ranging from high alpine peaks and northern boreal forest to rolling plateaus, deep canyons, and desert sage-steppe plains (Figure 2). This landscape is home to iconic North American wildlife and fish such as the grizzly bear and wild Pacific salmon; it includes some of the largest, most intact, and most protected areas outside of Alaska.

Landcover throughout the GNLCC is dominated by coniferous forests (44%) and shrublands (34%). Grasslands (9%), cropland (5%), barren land (3%), and water (2%) are other significant components (Figure 2). In the U.S. portion of the GNLCC, federal agencies (U.S. Forest Service, Bureau of Land Management, National Park Service, Bureau of Reclamation, and U.S. Fish and Wildlife Service) manage 53% of the land base; 37% is held privately; and states own 5%. In Canada, 94% of the landbase is Crown (public) land.

Landscape Stressors and Impacts

The overall ecological integrity of the Great Northern geography is impacted by landscape-level stressors. The three main stressors considered in this framework are climate change, invasive species, and land use change. The interaction of these stressors contributes to loss of habitat, fragmentation, changes in ecological function, and impacts on terrestrial and aquatic integrity. These stressors act at a scale that challenges the capacity and resources of partners.

Climate Change

The climate is changing, and the effects are seen in the Great Northern's valuable ecosystems and in the economies and communities that depend on plants, animals, and habitats. Warmer temperatures

and changing precipitation patterns are expected to cause more fires and more pest outbreaks such as the mountain pine beetle epidemic, while boreal forest are expected to move north into what is now tundra. Grasslands and shrublands are more likely to be invaded by non-native species, and wetlands may suffer losses from drier conditions. Climate change can exacerbate the impacts of habitat fragmentation and loss of ecosystem integrity caused by other landscape stressors.

Invasive Species

Invasive species are a threat to native biodiversity and have major implications for the conservation of public and private lands, and the production potential of working landscapes. Invading landscapes at alarming rates, invasive species are adversely impacting aquatic and terrestrial ecological function. Landscape effects have been documented from a range of invasive species including blister rust, cheatgrass, and aquatic invasive fish, macroinvertebrates, and pathogens such as whirling disease.

Land Use Change

From wildlands to working lands, the GNLCC landscape comprises a complex matrix of land uses reflecting diverse and changing economic reliance on natural resources. One major impact affecting natural resource management in Great Northern landscapes is energy development. This includes conventional development such as oil and gas, renewable energy development such as wind farms, and the infrastructure of transmission corridors and roads to support them. Land use decisions require specific kinds of science and information on key wildlife, ecological and landscape function, in order to minimize the loss of habitat connectivity and ecological function during development and operation of energy infrastructure. Increasing human populations in the regions instigate urban and exurban development and lead to increased human-use impacts on connectivity and ecological integrity.

Conservation Goals

To accomplish the vision of a landscape that sustains its diverse natural systems, the GNLCC intends to work toward ecological integrity at a landscape scale. Ecological integrity is defined as the ability of an ecological system to support and maintain a community of organisms that has the species composition, diversity, and functional organization comparable to those of natural habitats. Areas of highest ecological integrity have unfragmented natural landscapes, biotic and abiotic components well within the natural range of variability, and few impacts from invasive species. These areas are resilient to change, often contain large intact blocks of land, and sustain healthy and connected populations of fish, wildlife, and plants.

Landscape goals and interrelated, scaled conservation targets (ecosystem processes, ecosystems, and species) reflect the shared objectives of GNLCC partners.

Goals

This strategic framework maps hierarchically scaled interrelationships among conservation targets in order to allow partners working at all scales (local, regional, and national) to understand how their science and conservation efforts tier to the priorities of the GNLCC and contribute toward the over-arching goal of ecological integrity. These are described in greater detail in the following sections (see Figure 3).




Figure 3. Summary of the hierarchical relationships among conservation targets. [Note: additional species (elk, flammulated owl, black-backed woodpecker, flannelmouth sucker, blueheaded sucker, and roundtail chub) were also suggested, but have not been fully vetted, so are not included at this time.]

Goal 1: Maintain large, intact landscapes of naturally functioning terrestrial and aquatic community assemblages.

Large landscape conservation aims to ensure that large, intact areas connect and sustain diverse habitat types, and maintain ecological function and resilience in the context of existing stressors. The region's biodiversity exists in a mosaic of landscape elements including diverse habitat types across a variety of scales. Areas of refugia, connectivity, and smaller habitats are necessary to ensure resilience and the interactions among species and the ecological processes that support high-integrity systems.

Goal 2: Conserve a permeable landscape with connectivity across aquatic and terrestrial ecosystems, including species movement, genetic connectivity, migration, dispersal, life history, and biophysical processes (recognizing this is species dependent, and recognizing in some circumstances connectivity is not desired).

Habitats are likely to shift due to climate change and other stressors. In order to maintain desired connectivity (and limit connectivity in certain cases to minimize detrimental impacts), we need to understand the current level of landscape permeability, identify existing corridors, and anticipate how these might shift or change in condition in the future. Maintaining and restoring connectivity can help species adapt to climate change and mitigate other impacts of landscape stressors.

Goal 3: Maintain hydrologic regimes that support native or desirable aquatic plant and animal communities in still and moving water systems.

Hydrologic regimes (e.g., the timing, magnitude, and quality of water flow) are likely to shift due to climate change and other stressors. The ecological structure (e.g., configuration, connectivity, and size) and function of native or desirable riparian, aquatic, and wetland community assemblages provide habitat for aquatic-dependent species. Maintaining the functional matrix of aquatic and hydrologic systems is critical to maintain aquatic biodiversity, resilience, and ecosystem processes.

Goal 4: Promote landscape-scale disturbance regimes that operate within a future range of variability and sustain ecological integrity.

Disturbance regimes are shifting in time and space due to climate change (e.g., frequency and intensity of fire, spread of forest insects and disease, frequency and severity of drought and flooding). Risk assessments can help identify future conditions and the potential for extreme events, help identify vulnerable areas, and focus management efforts on areas with the greatest ecological contribution to the landscape. Maintaining ecosystem processes within the range of variability of disturbance can facilitate climate adaptation efforts.

Conservation Targets

The GNLCC has conceptually related broad conservation issues to three “ecotypic areas”—Columbia Basin, Rocky Mountain, and Sagebrush Steppe. These ecotypic areas cover large expanses within the GNLCC, have similar ecological systems with relatively consistent species compositions and configurations, and are facing common conservation issues. Our intent in identifying ecotypic areas is to connect local landscape partnerships working on similar conservation issues to broaden their positive affects to larger landscape scales.

Species were identified based on the documentation of their importance in relevant foundational documents, their representation of the ecotypic area, and their occurrence and distribution throughout the GNLCC landscape. Species were considered based on their role as: (Tier 1) umbrella species whose function and outcomes represent many other species; (Tier 2) ecological process indicator species

that have direct ties to critical ecological processes, which can indicate change or threshold effects; and (Tier 3) species of iconic or social importance, which serve as a flagship for landscape effects. Table 1 provides an example of how species are related to each goal.

Figure 4 is a conceptual model that demonstrates, through simple relationship arrows, the complexity of the interrelationships among these scaled conservation targets. Additional analysis will examine the species-species (trophic interaction), species-habitat, and species-landscape relationships. It is expected that these conceptual models and the information therein, will be refined by expert review and in the development of the GNLCC science strategy. In the meantime, these models provide an example of how specific ecological information can be used to elucidate complexity and inter-relatedness.



PHOTO CREDIT: iSTOCK

Table 1. An example of species relationships to ecological integrity goals¹.

	Columbia Basin	Rocky Mountain	Sage Steppe
Goal 1: Maintain large intact, resilient landscapes of naturally functional terrestrial and aquatic community assemblages.	T1, T3: Salmon	T1, T3: Wolverine Grizzly bear Canadian lynx	T1: Sage grouse
Goal 2: Conserve a permeable landscape with connectivity across aquatic and terrestrial ecosystems, including species movement, genetic connectivity, migration, dispersal, life history, and biophysical processes.	T3: Mule deer	T2: Wolverine Canadian lynx	T1: Sage grouse T2: Antelope Burrowing owl Pygmy rabbit T3: Mule deer
Goal 3: Maintain hydrologic regimes support native or desirable aquatic plant and animal communities in still and moving water systems.	T2: Salmon Steelhead Bull trout	T2: Bull trout Cutthroat trout Trumpeter swan	T2: Cutthroat trout
Goal 4: Promote landscape-scale disturbance regimes that operate within a future range of variability that sustains ecological integrity.	T2: Bull trout	T2: Wolverine Whitebark pine T3: Bull trout Cutthroat trout	T2: Sage grouse Burrowing owl

¹Species listed here and throughout the framework are examples based on input to date and may be refined or changed over time.




Figure 4. A conceptual model that demonstrates, through simple relationship arrows, an example of the complexity of the inter-relationships among scaled conservation targets.

Columbia Basin

The Columbia River drains a 259,000-square-mile basin that includes portions of several states of the interior western U.S. and British Columbia. The river is arguably the most significant environmental force and ecological lifeline in the Pacific Northwest and flows for more than 1,200 miles from the base of the Canadian Rockies to the Pacific Ocean. The Columbia River is a snow-charged river that seasonally fluctuates in volume with an annual average discharge of 160 million acre-feet of water, the highest volumes between April and September, and the lowest from December to February. Although people have lived along the river for more than 10,000 years, modern engineering in the 19th and 20th centuries has dramatically altered the hydrology and ecology of the Columbia.

Stressors and Impacts

The expected effects of climate change in the Columbia Basin include: earlier spring runoffs, greater peak flows, and reduced late-summer flows that will cause many smaller tributaries to become intermittent and reduce wetlands. Increased erosion from flood events and lower in-stream flow increase water temperature, which favors warm-water fish and a decreased survival of native cold-water fish. A better

understanding of how climate changes and water use effect hydrologic regimes is critical.

Human water development and use are a major ecological impact in the Columbia Basin. The invasion of exotic aquatic species create competition with native fish or damage natural habitats; and expected temperature increases are incompatible with native cold-water fish and may encourage the persistence and dominance of non-native species.

Conservation Targets

The riverine and riparian systems of the Columbia River, the Snake River, and their many tributaries define the local watersheds and biological diversity of the Columbia Basin. Critical landscape-scale ecological processes in the Columbia Basin include connectivity of aquatic systems and hydrologic function. Wetlands and the watershed uplands are important to ecological and hydrological function as they provide important habitat and maintain the health of the riparian and aquatic systems. To maintain these ecosystems and their functionality, a set of interrelated conservation targets have been identified for this ecotype (see Table 2).

Table 2. Summary of conservation targets for the Columbia Basin ecotypic area.

Ecosystem Processes	Habitats and Ecosystems	Species
aquatic connectivity	riverine	salmon
	riparian corridors	steelhead trout
	wetlands	bull trout
	watershed uplands	mule deer

For thousands of years, salmon thrived in the Columbia Basin. In less than 150 years, salmon have been driven to the brink of extinction due to barriers to their migration and degradation of habitat. Salmon are central to Native American cultures in the Columbia Basin, and are considered among the sacred “First Foods.” Extensive recovery efforts are underway. Steelhead trout are sea-run rainbow trout that, like salmon, return to their freshwater hatching grounds to spawn. They are impacted by the same threats as salmon and are also valued by Native American cultures. Bull trout have exacting niche-habitat demands, requiring water temperatures generally below 55°F (13°C), clean gravel beds, deep pools, complex cover such as snags and cut banks, and large systems of interconnected waterways to

accommodate spawning migrations. Thus, they favor the deep pools of cold lakes and large rivers, as well as high, cold mountain headwaters. Bull trout require clean, cold water for survival; are sensitive to habitat degradation; and have declined due to the introduction of non-native fish species.

Mule deer are common and widely distributed throughout the GNLCC region. The mule deer is significant as one of the traditional “First Foods” of Native American tribes and an economically important species to state-based wildlife management and conservation programs. They are an important species when considering the connections between the aquatic and riparian ecosystems with watershed uplands and are sensitive to the loss of sage-steppe habitat.



PHOTO CREDIT: National Park Service

Rocky Mountains

The Rocky Mountains are a dominant geomorphic feature of the GNLCC landscape. We consider mountain ecotypes in other parts of the Great Northern area, such as the Cascades of eastern Washington and Oregon as part of the Rocky Mountain ecotype. Generally, the mountain environment is divided into three main life zones: montane, subalpine, and alpine. Each provides a unique mixture of climate, landscape, and vegetation.

The Rocky Mountains provide habitat for a diverse variety of wildlife species, and most importantly, provide the connectivity and extent of habitat needed for wide-ranging, reclusive species such as grizzly bear, wolverine, and numerous other iconic and wide-ranging species.

Stressors and Impacts

The results of climate change expected in this ecotypic area include: glaciers retreating more rapidly, drought and fire regimes changing and becoming more frequent, invasive species and forest diseases spreading more rapidly, and streams generally warming. The land use conversion of this ecosystem to exurban residential development, energy development, and human infrastructure

is a major stressor that can result in fragmentation and a loss of connectivity. Invasive species impact both terrestrial and aquatic ecosystems degrading ecological integrity and threatening native species.

Conservation Targets

These Rocky Mountain ecosystems are unique to the GNLCC. Many wildlife populations in the Rocky Mountains rely on habitat connectivity and corridors to range widely based on the size of their home ranges, seasonal uses of habitats, and in some cases, specific migration routes. As climate change and other stressors impact the landscape, more information is needed to better understand how those ranges and corridors might shift and change in condition. The health of montane woodlands and other habitats are critical to maintaining ecological integrity. The modification of natural fire regimes, warming trends that have facilitated the increase of pine bark beetles, and drought stress have all contributed to a decline in forest health in the region. To maintain these ecosystems and their functionality, a set of interrelated conservation targets have been identified for this ecotype (see Table 3).

Table 3. Summary of conservation targets for Rocky Mountain ecotypic area.

Ecosystem Processes	Habitats and Ecosystems	Species
connectivity	alpine	grizzly bear
natural fire regimes	sub-alpine	wolverine
insects and forest pathogens	woodland	Canada lynx
	sage shrub/grasslands	whitebark pine
	alpine lakes	trumpeter swan
	rivers and riparian corridors	cutthroat trout
	wetlands	bull trout

The Rocky Mountains include diverse ecosystems. The mosaic of these diverse systems and their connectivity is critical to maintaining ecological integrity. Riparian systems support some of the highest biological diversity and crucial wildlife habitat in this mountainous region, as they help connect the many ecosystems. Alpine and montane waterways are sensitive and critical to maintain native fish species. Alpine ecosystems are particularly sensitive to the impacts of climate change, and these impacts may be more readily evident as we study ecological shifts.

Some species, such as the grizzly bear, wolverine, and lynx, are wide-ranging species iconic to this region. Grizzly bear are more general in their habitat requirements, but whose key habitats provide an umbrella for many others; whereas a species like wolverine is dependent on conditions related to climate and snowpack and can serve as an indicator for certain landscape- and climate-related processes.

Whitebark pine is typically the highest-elevation pine tree of the Rocky Mountains, and is an important food source for many species—from grizzly bear to seed-eating birds and small mammals. Non-native blister rust, a tree pathogen, and

infestations of native pine beetles have dramatically impacted the populations of whitebark pine in this region. As a result, whitebark pine is declining throughout its range and was recently listed as “warranted” but precluded for status as an endangered species.

Although the Rocky Mountain population of trumpeter swan is healthy, some locations in the heart of its range (the greater Yellowstone region) are experiencing a decrease in number of birds and reproductive success.

Native fish species in the Rocky Mountains are threatened by changing hydrologic cycles (and warming water temperatures) and invasive nonnative species. Bull trout have exacting niche-habitat demands, requiring water temperatures generally below 55°F (13°C), clean gravel beds, deep pools, complex cover such as snags and cut banks, and large systems of interconnected waterways to accommodate spawning migrations. Thus, they favor the deep pools of cold lakes and large rivers, as well as high, cold mountain headwaters. Bull trout are sensitive to disturbance. Cutthroat trout have evolved into a number of subspecies, several of which are native to the GNLCC.

PHOTO CREDIT: National Park Service



Sagebrush Steppe

Sagebrush steppe ecosystems are some of the most threatened in North America and represent some of the most unique shrubland systems in the world. These semi-arid shrub and grassland systems are typical of broad stretches of lowlands in the Intermountain West, and occur in the Columbia Basin as well as montane valleys of Wyoming and Montana. The sagebrush steppe region is characterized by sagebrush and perennial bunchgrasses in semi-arid uplands occurring between 1,900 and 6,900 feet in elevation.

Stressors and Impacts

The results of climate change expected in the sage-steppe include an increase in temperature and shift in precipitation from summer to winter. This is predicted to favor pinyon pine-juniper woodland expansion and to create a positive feedback loop of annual grass establishment and greater fire frequency.

The land use conversion of this ecosystem to agriculture, energy development, residential development, and human infrastructure (e.g., transmission corridors, water development, and fencing, among others) is a major stressor. Invasive species such as cheatgrass can become established due to soil disturbance, shrub mortality,

fire, and the timing of rainfall (i.e., winter/early spring rains encourage the growth of annual exotics over native bunchgrasses, which are dependent on summer rains). Once these exotic annual grasses become established, they alter the fire regime and encourage more frequent, intense, and extensive fires, which in turn favor annual grasses over shrubs and native forbs.

Conservation Targets

The semi-arid, sage-dominated shrub and grasslands are the primary focal ecosystems for this ecotype. It is recognized that transition zones, riparian areas, pothole lakes, and other aquatic ecosystems are critical to maintaining ecosystem function and various ecosystem components. Sage-steppe ecosystems are threatened by habitat conversion, encroachment, and fragmentation by roads, land cover conversions, and other infrastructure. Connectivity is needed to sustain ecosystem integrity. The invasion of exotic plants and grasses has altered natural fire regimes in sagebrush/grassland ecosystems in a way that is advantageous to these invasive species. To maintain these ecosystems and their functionality, a set of interrelated conservation targets have been identified for this ecotype (see Table 4).

Table 4. Summary of conservation targets for Sage-steppe ecotypic area.

Ecosystem Processes	Habitats and Ecosystems	Species
connectivity	sage shrub/grassland	greater sage grouse
natural fire regimes	riparian corridors	pygmy rabbit
	pothole lakes	burrowing owl
		pronghorn antelope
		mule deer
		cutthroat trout

The greater sage grouse is being considered for protection under the Endangered Species Act. Evidence suggests that habitat fragmentation and destruction across much of the species' range has contributed to significant population declines over the past century. Other sagebrush obligate species include the pygmy rabbit and burrowing owl. Burrowing owls are commonly found in sage-steppe areas where shrub cover is less-dense. However, they remain susceptible to negative impacts, especially those which expose them to increased predation risk.

Pronghorn are relatively common species in the sage-dominated arid grasslands, however, they are North America's most wide-ranging migratory mammal and many migration corridors are threatened

by landcover and land-use changes that block traditional migration routes. Mule deer are also common and widely distributed throughout the GNLCC region, but significant as one of the traditional First Foods of Native American tribes and an economically important species to state-based wildlife management and conservation programs. Both pronghorn and mule deer need large areas of diverse habitat and connected landscape.

Native fish species are also important in the sagebrush steppe region. They are threatened by changing hydrologic cycles (including water quantity and quality impacted by energy development) and invasive nonnative species. For example, cutthroat trout have cold-water requirements and have been identified as high priority for conservation in the region.



PHOTO CREDIT: National Park Service

Next Steps

The Great Northern LCC is a network of partners working toward common goals. This strategic conservation framework is intended to provide the landscape-level institutional and conceptual frameworks to facilitate cooperative and responsive adaptive management in the face of three common large scale stressors: climate change, land development, and invasive species.

The Great Northern LCC has identified an initial set of collective conservation targets that have been developed through a preliminary participatory process among federal, state, and tribal land management agencies and non-governmental science stakeholders. The purpose of these conservation targets is to frame and focus the work of the Great Northern LCC and address large-scale landscape issues critical to this region. The strategies identified by the GNLCC partners incorporate the scale-appropriate conservation actions of existing efforts and at the same time, the larger goals identified for the entire breadth of the GNLCC.

The next steps for implementing the GNLCC strategy include:

- Building organizational capacity by supporting the partner forums to work collaboratively and effectively to effect landscape conservation at the broadest scale.
- Integrating the Cascadia Forum, a self-directed group supported by both the GNLCC and the North Pacific LCC, into GNLCC partner forum discussions.

- Aligning and sharing existing information and conducting a gap analysis to determine how and where GNLCC efforts are most strategically invested.
- Developing a science strategy based on the gap analysis to guide strategic science investment.
- Partnering with inventory and monitoring programs to collect and share information and cross-walk data across agencies and organizations.
- Identifying resource managers' needs for information and present that information in a useful, accessible way and developing tools to disseminate information and inform management decisions.
- Creating a communications strategy to keep the GNLCC network of partners engaged and informed.
- Identifying opportunities to focus and leverage resources on priority issues.
- Linking conservation to sustainable communities through socioeconomic analysis.

In pursuing these steps, the GNLCC and its partners seek to manage the region in an adaptive fashion by incorporating new information and adjusting strategies accordingly.

Acknowledgements

We are grateful for the contributions of partners and advisory team members who helped shape this document. We also thank Craig Groves at The Nature Conservancy who provided valuable guidance throughout this process.

Tyler Abbot, U. S. Fish and Wildlife Service
Dan Blake, U. S. Fish and Wildlife Service
Zach Bowen, USGS, Wyoming Landscape Conservation Initiative
Anne Carlson, The Wilderness Society
Molly Cross, Wildlife Conservation Society
Frank D'Erchia, USGS, Wyoming Landscape Conservation Initiative
Bill Dolan, Alberta Parks
John Emmerich, Wyoming Game and Fish Department
Wendy Francis, Yellowstone to Yukon Conservation Initiative
Jon Haber, U. S. Forest Service
Beth Hahn, U. S. Forest Service
Sonia Hall, The Nature Conservancy, Arid Lands Initiative
Virginia Kelly, Greater Yellowstone Coordinating Committee
Nancy Lee, USGS, Northwest Climate Science Center
Kaaren Lewis, British Columbia Ministry of the Environment
Steve Lozar, Confederated Salish Kootenai Tribe
Madeline Maley, British Columbia Ministry of Forests and Range
Mary Manning, U. S. Forest Service
Jim Morrison, USFS, Northern Forest Regional Climate Coordinator
Jeff Morissette, U. S. Geological Survey
John Pierce, Washington Department of Fish and Wildlife
Tasha Sargent, Environment Canada, Canadian Wildlife Service
Carl Scheeler, Confederated Tribes of the Umatilla Indian Reservation
Gregg Servheen, Idaho Department of Fish and Game
Erin Sexton, Crown Managers Partnership
Brad Shepard, Wildlife Conservation Society
Kathy Tonnessen, NPS, Rocky Mountain Cooperative Ecosystems Studies Unit
Steve Torbitt, U.S. Fish and Wildlife Service
Rich Torquemada, Bureau of Land Management
Josh Vest, USFWS, Intermountain Joint Venture
Steve Waste, U. S. Geological Survey
Michael Whitfield, Heart of the Rockies
Styphen Zylstra, U. S. Fish and Wildlife Service

PHOTO CREDIT: Gary Tabor





Great Northern LCC Coordinators

Yvette Converse, Coordinator
406.994.7486
Yvette_Converse@fws.gov

Tom Olliff, Coordinator
406.994.7920
Tom_Olliff@nps.gov

Sean Finn, Science Coordinator
208.426.2697
Sean_Finn@fws.gov

Rick Sojda, Science Coordinator
406.994.6556
sojda@usgs.gov

January 2013

greatnorthernlcc.org

PHOTO CREDIT: Gary Tabor